PAC Learning

Consider a class C of possible target concepts defined over a set of instances X of length n, and a learner L using hypothesis space H.

> *Definition:* C is **PAC-learnable** by L using H if for all $c \in C$, distributions $\mathcal{D}$ over X, ϵ such that $0 < \epsilon < 1/2$, and δ such that $0 < \delta < 1/2$,
>
> learner L will with probability at least $(1 - \delta)$ output a hypothesis $h \in H$ such that $error_{\mathcal{D}}(h) \leq \epsilon$, in time that is polynomial in $1/\epsilon$, $1/\delta$, n and $size(c)$.

Shattering a Set of Instances

Definition: a **dichotomy** of a set S is a partition of S into two disjoint subsets.

Definition: a set of instances S is **shattered** by hypothesis space H if and only if for every dichotomy of S there exists some hypothesis in H consistent with this dichotomy.

Three Instances Shattered

Instance space X



The Vapnik-Chervonenkis Dimension

Definition: The **Vapnik-Chervonenkis dimension**, $VC(H)$, of hypothesis space H defined over instance space X is the size of the largest finite subset of X shattered by H. If arbitrarily large finite sets of X can be shattered by H, then $VC(H) \equiv \infty$.

VC Dim. of Linear Decision Surfaces

(a) (b)

Sample Complexity from VC Dimension

How many randomly drawn examples suffice to ϵ-exhaust $VS_{H,D}$ with probability at least $(1-\delta)$?

$$m \geq \frac{1}{\epsilon}(4\log_2(2/\delta) + 8VC(H)\log_2(13/\epsilon))$$