


Surfaces of Revolution

**Brian Curless
CSEP 557
Fall 2016**


Surfaces of revolution



Idea: rotate a 2D **profile curve** around an axis.

What kinds of shapes can you model this way?

Constructing surfaces of revolution



Given: A curve $C(v)$ in the xy -plane:

$$C(v) = \begin{bmatrix} C_x(v) \\ C_y(v) \\ 0 \\ 1 \end{bmatrix}$$


Let $R_y(\theta)$ be a rotation about the y -axis.

Find: A surface $S(u, v)$ which is $C(v)$ rotated about the y -axis, where $u, v \in [0, 1]$.

Solution: $S(u, v) = R_y(2\pi u)C(v)$

Constructing surfaces of revolution

We can sample in u and v to get a grid of points over the surface.



Suppose we sample:

- ♦ in v , to give $C[j]$ where $j \in [0..M-1]$
- ♦ in u , to give rotation angle $\theta[i] = 2\pi i/N$ where $i \in [0..N]$

We can now write the surface as:


$$S[i,j] = R_y\left(2\pi \frac{i}{N}\right) C[j]$$

How would we turn this into a mesh of triangles?
How do we assign per-vertex normals?

Surface normals


Now that we describe the surface as a triangle mesh, we need to provide surface normals. As we'll see later, these normals are important for drawing and shading the surface (i.e., for "rendering").

One approach is to compute the normal to each triangle. How do we compute these normals?




For surfaces of revolution, we can get better-looking results by analytically computing the normal at each vertex...

Tangent vectors, tangent planes, and normals



$$t \approx Q - P$$

$$t = \lim_{Q \rightarrow P} \frac{Q - P}{\|Q - P\|}$$




$$t_1 \approx Q - P$$

$$t_2 \approx R - P$$

$$N = \frac{t_2 \times t_1}{\|t_2 \times t_1\|}$$

Normals on a surface of revolution



We can compute tangents to the curve points in the x-y plane:

$$T_1[0, j] \approx c[j+1] - c[j]$$

$$T_2[0, j] = [0 \ 0 \ 1]^T$$

to get the normal in that plane:

$$N[0, j] = T_1[0, j] \times T_2[0, j] \leftarrow \text{normalize this}$$

and then rotate it around:

$$N[i, j] = R_y\left(2\pi \frac{i}{n}\right) N[0, j]$$

Triangle meshes

How should we generally represent triangle meshes?

Δ by Δ list

V_1, N_1


V_5, N_5

V_6, N_6

V_1, N_1

V_6, N_6

V_2, N_2



Vertex list + Δ index list

V_1, N_1

V_2, N_2

V_3, N_3

V_4, N_4

V_5, N_5

V_6, N_6

1, 5, 6

1, 6, 2

Do this

Summary

What to take away from this lecture:

- ◆ All the names in boldface.
- ◆ How to compute a surface of revolution given a profile curve.
- ◆ How to represent a surface of revolution as a triangle mesh.
- ◆ How to compute per-vertex normals for a surface of revolution.