

# Affine transformations

# Reading

Optional reading:

- ♦ Angel and Shreiner: 3.1, 3.7-3.11
- ♦ Marschner and Shirley: 2.3, 2.4.1-2.4.4, 6.1.1-6.1.4, 6.2.1, 6.3

Further reading:

- ♦ Angel, the rest of Chapter 3
- ♦ Foley, et al, Chapter 5.1-5.5.
- ♦ David F. Rogers and J. Alan Adams, *Mathematical Elements for Computer Graphics*, 2<sup>nd</sup> Ed., McGraw-Hill, New York, 1990, Chapter 2.

# Geometric transformations

Geometric transformations will map points in one space to points in another:  $\underline{(x', y', z')} = \underline{f}(x, y, z)$ .

These transformations can be very simple, such as scaling each coordinate, or complex, such as non-linear twists and bends.

We'll focus on transformations that can be represented easily with matrix operations.


# Vector representation

We can represent a **point**,  $\mathbf{p} = (x, y)$ , in the plane or  $\mathbf{p} = (x, y, z)$  in 3D space:

- ♦ as column vectors  $\begin{bmatrix} x \\ y \end{bmatrix}$   $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$

- ♦ as row vectors  $\begin{bmatrix} x & y \end{bmatrix}$   $\begin{bmatrix} x & y & z \end{bmatrix}$


# Canonical axes




right hand rule

$$v = x\hat{x} + y\hat{y} + z\hat{z}$$
$$= x \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + y \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + z \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$



# Vector length and dot products



$$u = \begin{bmatrix} u_x \\ u_y \\ u_z \end{bmatrix} \quad v = \begin{bmatrix} v_x \\ v_y \\ v_z \end{bmatrix}$$

$$\|v\| = \sqrt{v_x^2 + v_y^2 + v_z^2}$$

$$\hat{v} = \frac{v}{\|v\|} \Rightarrow \|\hat{v}\| = 1$$

$$u \cdot v = u_x v_x + u_y v_y + u_z v_z$$

$$u \cdot v \stackrel{?}{=} u^T v$$

$$u \cdot v \stackrel{?}{=} v \cdot u \quad \checkmark$$

$$\stackrel{?}{=} \begin{bmatrix} u_x & u_y & u_z \end{bmatrix} \begin{bmatrix} v_x \\ v_y \\ v_z \end{bmatrix}$$

$$v \cdot v = \|v\|^2$$

$$u \cdot v = \|u\| \|v\| \cos \theta$$

$$u \cdot v = 0 \Rightarrow u \perp v \quad \text{perpendicular, orthogonal}$$

$\|u\|, \|v\| \neq 0$

$$\hat{u} \cdot \hat{v} = \cos \theta$$

$$(\|\hat{u}\| = \|\hat{v}\| = 1)$$

## Representation, cont.

We can represent a **2-D transformation**  $M$  by a matrix

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

If  $\mathbf{p}$  is a column vector,  $M$  goes on the left:

$$\mathbf{p}' = M\mathbf{p}$$

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} ax+by \\ cx+dy \end{bmatrix}$$

If  $\mathbf{p}$  is a row vector,  $M^T$  goes on the right:

$$\mathbf{p}' = \mathbf{p}M^T$$

$$\begin{bmatrix} x' & y' \end{bmatrix} = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} a & c \\ b & d \end{bmatrix} = \begin{bmatrix} ax+by & cx+dy \end{bmatrix}$$

We will use **column vectors**.

$$(AB)^T = B^T A^T$$

$$(AB)^{-1} = B^{-1} A^{-1}$$

$$(AB)^{-1}(AB) = I$$

$$(AB)^{-1}AB = I$$

$$(AB)^{-1}A = B^{-1}$$

$$(AB)^{-1} = B^{-1}A^{-1}$$



# Two-dimensional transformations

Here's all you get with a 2 x 2 transformation matrix  $M$  :

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

So:

$$x' = ax + by$$

$$y' = cx + dy$$

We will develop some intimacy with the elements  $a$ ,  $b$ ,  $c$ ,  $d$ ...



# Identity

Suppose we choose  $a = d = 1, b = c = 0$ :

- ◆ Gives the **identity** matrix:

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

- ◆ Doesn't move the points at all

$$x' = x$$

$$y' = y$$

# Scaling

Suppose we set  $b = c = 0$ , but let  $a$  and  $d$  take on any *positive* value:



- ◆ Gives a **scaling** matrix:

$$\begin{bmatrix} a & 0 \\ 0 & d \end{bmatrix}$$


- ◆ Provides **differential (non-uniform) scaling** in  $x$  and  $y$ :

$$x' = ax$$

$$y' = dy$$



$$\begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ 2y \end{bmatrix}$$



$$\begin{bmatrix} 1/2 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{1}{2}x \\ 2y \end{bmatrix}$$



## Mirror / reflection

Suppose we keep  $b = c = 0$ , but let either  $a$  or  $d$  go negative.

Examples:

$$\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -x \\ y \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -x \\ -y \end{bmatrix}$$



rotation by  $180^\circ$

# Shear

Now let's leave  $a = d = 1$  and experiment with  $b \dots$



The matrix

$$\begin{bmatrix} 1 & b \\ 0 & 1 \end{bmatrix}$$

gives:

$$x' = x + by$$

$$y' = y$$



$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x+y \\ y \end{bmatrix}$$





# Effect on unit square

Let's see how a general 2 x 2 transformation  $M$  affects the unit square:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} \mathbf{q} & \mathbf{r} & \mathbf{s} & \mathbf{t} \end{bmatrix} = \begin{bmatrix} \mathbf{q}' & \mathbf{r}' & \mathbf{s}' & \mathbf{t}' \end{bmatrix}$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 0 & a & a+b & b \\ 0 & c & c+d & d \end{bmatrix}$$

$\uparrow \quad \uparrow \quad \uparrow$



## Effect on unit square, cont.



Observe:

- ♦ Origin invariant under  $M$
- ♦  $M$  can be determined just by knowing how the corners  $(1,0)$  and  $(0,1)$  are mapped
- ♦  $a$  and  $d$  give  $x$ - and  $y$ -scaling
- ♦  $b$  and  $c$  give  $x$ - and  $y$ -shearing

# Rotation

soh cah toa

From our observations of the effect on the unit square, it should be easy to write down a matrix for "rotation about the origin":



$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} \cos\theta \\ \sin\theta \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -\sin\theta \\ \cos\theta \end{bmatrix}$$

Thus,

$$M = R(\theta) = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$$

# Limitations of the 2 x 2 matrix

A 2 x 2 linear transformation matrix allows

- ◆ Scaling
- ◆ Rotation
- ◆ Reflection
- ◆ Shearing

**Q:** What important operation does that leave out?

*translation*



# Homogeneous coordinates



Idea is to lift the problem up into 3-space, adding a third component to every point:

$$\begin{bmatrix} x \\ y \end{bmatrix} \rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Adding the third "w" component puts us in **homogenous coordinates**.

And then transform with a 3 x 3 matrix:

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = T(\mathbf{t}) \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix}$$



... gives **translation!**

# Anatomy of an affine matrix

The addition of translation to linear transformations gives us **affine transformations**.

In matrix form, 2D affine transformations always look like this:

$$M = \begin{bmatrix} a & b & t_x \\ c & d & t_y \\ 0 & 0 & 1 \end{bmatrix} = \left[ \begin{array}{cc|c} A & & \mathbf{t} \\ \hline 0 & 0 & 1 \end{array} \right]$$

$$M \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} ax + by + t_x \\ cx + dy + t_y \\ 1 \end{bmatrix}$$

2D affine transformations always have a bottom row of  $[0 \ 0 \ 1]$ .

An "affine point" is a "linear point" with an added  $w$ -coordinate which is always 1:

$$\mathbf{p}_{\text{aff}} = \begin{bmatrix} \mathbf{p}_{\text{lin}} \\ 1 \end{bmatrix} = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Applying an affine transformation gives another affine point:

$$M \mathbf{p}_{\text{aff}} = \begin{bmatrix} A \mathbf{p}_{\text{lin}} + \mathbf{t} \\ 1 \end{bmatrix}$$


# Rotation about arbitrary points

Until now, we have only considered rotation about the origin.

With homogeneous coordinates, you can specify a rotation by  $\beta$ , about any point  $\mathbf{q} = [q_x \ q_y]^T$  with a matrix.

Let's do this with rotation and translation matrices of the form  $R(\theta)$  and  $T(\mathbf{t})$ , respectively.

$$R(\theta) = \begin{bmatrix} \cos & -\sin & 0 \\ \sin & \cos & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$T(\mathbf{t}) = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}$$



$$M = T(\mathbf{q})R(\beta)T(-\mathbf{q})$$

1. Translate  $\mathbf{q}$  to origin
2. Rotate
3. Translate back

# Points and vectors

Vectors have an additional coordinate of  $w = 0$ . Thus, a change of origin has no effect on vectors.

**Q:** What happens if we multiply a vector by an affine matrix?

$$\begin{bmatrix} a & b & t_x \\ c & d & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} v_x \\ v_y \\ 0 \end{bmatrix} = \begin{bmatrix} a v_x + b v_y \\ c v_x + d v_y \\ 0 \end{bmatrix}$$

$$\alpha \begin{bmatrix} A_x \\ A_y \\ 1 \end{bmatrix} + \beta \begin{bmatrix} B_x \\ B_y \\ 1 \end{bmatrix} = \begin{bmatrix} \alpha A_x + \beta B_x \\ \alpha A_y + \beta B_y \\ \alpha + \beta \end{bmatrix}$$

These representations reflect some of the rules of affine operations on points and vectors:

- vector + vector → vector
- scalar · vector → vector
- point - point → vector
- point + vector → point
- point + point → chaos


- $\alpha A + \beta B$
- $\alpha + \beta = 0 \Rightarrow$  vector
- $\alpha + \beta = 1 \Rightarrow$  point
- else  $\Rightarrow$  chaos

- scalar · vector + scalar · vector → vector
- scalar · point + scalar · point → it depends!

One useful combination of affine operations is:

$$P(t) = P_0 + t\mathbf{u}$$

$t \in (\infty, \infty) \Rightarrow$  line  
 $t \in [0, \infty) \Rightarrow$  half-line ray





**Q:** What does this describe?

# Basic 3-D transformations: scaling

Some of the 3-D transformations are just like the 2-D ones.



For example, scaling:

$$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 & 0 \\ 0 & s_y & 0 & 0 \\ 0 & 0 & s_z & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$$



# Translation in 3D

$$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & t_x \\ 0 & 1 & 0 & t_y \\ 0 & 0 & 1 & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$$



## Rotation in 3D (cont'd)


These are the rotations about the canonical

axes:

$$R_x(\alpha) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \alpha & -\sin \alpha & 0 \\ 0 & \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$R_y(\beta) = \begin{bmatrix} \cos \beta & 0 & \sin \beta & 0 \\ 0 & 1 & 0 & 0 \\ -\sin \beta & 0 & \cos \beta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$R_z(\gamma) = \begin{bmatrix} \cos \gamma & -\sin \gamma & 0 & 0 \\ \sin \gamma & \cos \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$



Use right hand rule


A general rotation can be specified in terms of a product of these three matrices. How else might you specify a rotation?

quaternion

# Shearing in 3D

Shearing is also more complicated. Here is one example:

$$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & b & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$$






We call this a shear with respect to the x-z plane.



# Properties of affine transformations

Here are some useful properties of affine transformations:

- ◆ Lines map to lines
- ◆ Parallel lines remain parallel
- ◆ Midpoints map to midpoints (in fact, ratios are always preserved)



$$\text{ratio} = \frac{\|pq\|}{\|qr\|} = \frac{s}{t} = \frac{\|p'q'\|}{\|q'r'\|}$$

# Summary

What to take away from this lecture:

- ◆ All the names in boldface.
- ◆ How points and transformations are represented.
- ◆ How to compute lengths, dot products, and cross products of vectors, and what their geometrical meanings are.
- ◆ What all the elements of a  $2 \times 2$  transformation matrix do and how these generalize to  $3 \times 3$  transformations.
- ◆ What homogeneous coordinates are and how they work for affine transformations.
- ◆ How to concatenate transformations.
- ◆ The mathematical properties of affine transformations.