


CSE 311: Foundations of Computing

Lecture 14: Induction



Mathematical Induction

Method for proving statements about all natural numbers

- A new logical inference rule!
 - It only applies over the natural numbers
 - The idea is to **use** the special structure of the naturals to prove things more easily
- Particularly useful for reasoning about programs!
`for (int i=0; i < n; n++) { ... }`
 - Show $P(i)$ holds after i times through the loop

Prove $\forall a, b, m > 0 \forall k \in \mathbb{N} ((a \equiv b \pmod{m}) \rightarrow (a^k \equiv b^k \pmod{m}))$

Let $a, b, m > 0$ be arbitrary. Let $k \in \mathbb{N}$ be arbitrary.

Suppose that $a \equiv b \pmod{m}$.

We know that by multiplying congruences we get

$$(a \equiv b \pmod{m} \wedge a \equiv b \pmod{m}) \rightarrow a^2 \equiv b^2 \pmod{m}$$

Then, repeating this many times, we have:

$$(a^2 \equiv b^2 \pmod{m} \wedge a \equiv b \pmod{m}) \rightarrow a^3 \equiv b^3 \pmod{m}$$

$$(a^3 \equiv b^3 \pmod{m} \wedge a \equiv b \pmod{m}) \rightarrow a^4 \equiv b^4 \pmod{m}$$

...

$$(a^{k-1} \equiv b^{k-1} \pmod{m} \wedge a \equiv b \pmod{m}) \rightarrow a^k \equiv b^k \pmod{m}$$

The “...” is a problem! We don’t have a proof rule that allows us to say “do this over and over”.


But there such a property of the natural numbers!

Domain: Natural Numbers

$$\begin{array}{l} \textcircled{1} P(0) \\ \textcircled{2} \forall k (P(k) \rightarrow P(k + 1)) \\ \hline \therefore \forall n P(n) \end{array}$$

Induction Is A Rule of Inference

Domain: Natural Numbers


$$\frac{P(0) \quad \forall k (P(k) \rightarrow P(k + 1))}{\therefore \forall n P(n)}$$


How do the givens prove $P(5)$?

Induction Is A Rule of Inference

Domain: Natural Numbers

$$\frac{P(0) \quad \forall k (P(k) \rightarrow P(k + 1))}{\therefore \forall n P(n)}$$

How do the givens prove P(5)?



First, we have **P(0)**.

Since $P(n) \rightarrow P(n+1)$ for all n , we have **P(0)** \rightarrow **P(1)**.

Since **P(0)** is true and **P(0)** \rightarrow **P(1)**, by Modus Ponens, **P(1)** is true.

Since $P(n) \rightarrow P(n+1)$ for all n , we have **P(1)** \rightarrow **P(2)**.

Since **P(1)** is true and **P(1)** \rightarrow **P(2)**, by Modus Ponens, **P(2)** is true.


...


Using The Induction Rule In A Formal Proof

$$\begin{array}{c} P(0) \\ \forall k (P(k) \rightarrow P(k + 1)) \\ \hline \therefore \forall n P(n) \end{array}$$

Using The Induction Rule In A Formal Proof

$$\begin{array}{l} \textcircled{1} P(0) \\ \textcircled{2} \forall k (P(k) \rightarrow P(k + 1)) \\ \hline \therefore \forall n P(n) \end{array}$$

 1. $P(0)$

 4. $\forall k (P(k) \rightarrow P(k+1))$
5. $\forall n P(n)$

Induction: 1, 4

Using The Induction Rule In A Formal Proof


$$\frac{P(0) \quad \forall k (P(k) \rightarrow P(k + 1))}{\therefore \forall n P(n)}$$

1. $P(0)$
2. Let k be an arbitrary integer ≥ 0
3. $P(k) \rightarrow P(k+1)$
4. $\forall k (P(k) \rightarrow P(k+1))$ Intro \forall : 2, 3
5. $\forall n P(n)$ Induction: 1, 4

Using The Induction Rule In A Formal Proof

$$\frac{P(0) \quad \forall k (P(k) \rightarrow P(k + 1))}{\therefore \forall n P(n)}$$

$\forall x,$
 $x \geq 0$
 $\rightarrow P(x)$

1. $P(0)$
2. Let k be an arbitrary integer ≥ 0 
 - 3.1. $P(k)$ Assumption
 - 3.2. ...
 - 3.3. $P(k+1)$
3. $P(k) \rightarrow P(k+1)$ Direct Proof Rule
4. $\forall k (P(k) \rightarrow P(k+1))$ Intro \forall : 2, 3
5. $\forall n P(n)$ Induction: 1, 4

Translating to an English Proof

$$P(0)$$
$$\forall k (P(k) \rightarrow P(k + 1))$$

$$\therefore \forall n P(n)$$

1. Prove $P(0)$

Base Case

2. Let k be an arbitrary integer ≥ 0

Inductive Hypothesis

3.1. Suppose that $P(k)$ is true

3.2. ...

Inductive Step

3.3. Prove $P(k+1)$ is true

3. $P(k) \rightarrow P(k+1)$

Direct Proof Rule

4. $\forall k (P(k) \rightarrow P(k+1))$

Intro \forall : 2, 3

5. $\forall n P(n)$

Induction: 1, 4

Conclusion

Translating to an English Proof

1. Prove $P(0)$	Base Case	
2. Let k be an arbitrary integer ≥ 0		Inductive Hypothesis
3.1. Assume that $P(k)$ is true		
3.2. ...		Inductive Step
3.3. Prove $P(k+1)$ is true		
3. $P(k) \rightarrow P(k+1)$		Direct Proof Rule
4. $\forall k (P(k) \rightarrow P(k+1))$		Intro \forall : 2, 3
5. $\forall n P(n)$		Induction: 1, 4
		Conclusion

Induction English Proof Template

[...Define $P(n)$...]

We will show that $P(n)$ is true for every $n \in \mathbb{N}$ by Induction.

Base Case: *[...proof of $P(0)$ here...]*

Induction Hypothesis:

Suppose that $P(k)$ is true for an arbitrary $k \in \mathbb{N}$.

Induction Step:

[...proof of $P(k + 1)$ here...]

*The proof of $P(k + 1)$ **must** invoke the IH somewhere.*

So, the claim is true by induction.

Inductive Proofs In 5 Easy Steps

Proof:

1. “Let $P(n)$ be... . We will show that $P(n)$ is true for every $n \geq 0$ by Induction.”

2. “Base Case:” Prove $P(0)$

3. “Inductive Hypothesis:

Suppose $P(k)$ is true for an arbitrary integer $k \geq 0$ ”

4. “Inductive Step:” Prove that $P(k + 1)$ is true.

Use the goal to figure out what you need.

Make sure you are using I.H. and point out where you are using it. (Don't assume $P(k + 1)$!!)

5. “Conclusion: Result follows by induction”

What is $1 + 2 + 4 + \dots + 2^n$?

- $1 = 1$
- $1 + 2 = 3$
- $1 + 2 + 4 = 7$
- $1 + 2 + 4 + 8 = 15$
- $1 + 2 + 4 + 8 + 16 = 31$

It sure looks like this sum is $\underbrace{2^{n+1} - 1}$

How can we prove it?

We could prove it for $n = 1, n = 2, n = 3, \dots$ but that would literally take forever.

Good that we have induction!

Prove $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$

Prove $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$

- 1. Let $P(n)$ be " $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ". We will show $P(n)$ is true for all natural numbers by induction.**

Prove $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$

- 1. Let $P(n)$ be " $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ". We will show $P(n)$ is true for all natural numbers by induction.**
- 2. Base Case ($n=0$): $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$ so $P(0)$ is true.**

Prove $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$

- 1. Let $P(n)$ be “ $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ”. We will show $P(n)$ is true for all natural numbers by induction.**
- 2. Base Case ($n=0$): $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$ so $P(0)$ is true.**
- 3. Induction Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 0$, i.e., that $2^0 + 2^1 + \dots + 2^k = 2^{k+1} - 1$.**

Prove $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$

- 1. Let $P(n)$ be “ $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ”. We will show $P(n)$ is true for all natural numbers by induction.**
- 2. Base Case ($n=0$): $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$ so $P(0)$ is true.**
- 3. Induction Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 0$, i.e., that $2^0 + 2^1 + \dots + 2^k = 2^{k+1} - 1$.**
- 4. Induction Step:**

Goal: Show $P(k+1)$, i.e. show $2^0 + 2^1 + \dots + 2^k + 2^{k+1} = 2^{k+2} - 1$

Prove $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$

1. Let $P(n)$ be " $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ". We will show $P(n)$ is true for all natural numbers by induction.
2. Base Case ($n=0$): $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$ so $P(0)$ is true.
3. Induction Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 0$, i.e., that $2^0 + 2^1 + \dots + 2^k = 2^{k+1} - 1$.
4. Induction Step:

$$2^0 + 2^1 + \dots + 2^k = 2^{k+1} - 1 \quad \text{by IH} \quad \leftarrow$$

Adding 2^{k+1} to both sides, we get:

$$2^0 + 2^1 + \dots + 2^k + 2^{k+1} = 2^{k+1} + 2^{k+1} - 1$$

Note that $2^{k+1} + 2^{k+1} = 2(2^{k+1}) = 2^{k+2}$.

So, we have $2^0 + 2^1 + \dots + 2^k + 2^{k+1} = 2^{k+2} - 1$, which is exactly $P(k+1)$.

Prove $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$

1. Let $P(n)$ be " $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ". We will show $P(n)$ is true for all natural numbers by induction.
2. Base Case ($n=0$): $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$ so $P(0)$ is true.
3. Induction Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 0$, i.e., that $2^0 + 2^1 + \dots + 2^k = 2^{k+1} - 1$.
4. Induction Step:

We can calculate

$$\begin{aligned} \underline{2^0 + 2^1 + \dots + 2^k + 2^{k+1}} &= \overbrace{(2^0 + 2^1 + \dots + 2^k)} + 2^{k+1} \\ &= (2^{k+1} - 1) + 2^{k+1} && \text{by the IH} \\ &= 2(2^{k+1}) - 1 \\ &= 2^{k+2} - 1, \end{aligned}$$

which is exactly $P(k+1)$.

Alternative way of writing the inductive step

Prove $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$ 

1. Let $P(n)$ be " $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ". We will show $P(n)$ is true for all natural numbers by induction.
2. Base Case ($n=0$): $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$ so $P(0)$ is true.
3. Induction Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 0$, i.e., that $2^0 + 2^1 + \dots + 2^k = 2^{k+1} - 1$.
4. Induction Step:

We can calculate

$$\begin{aligned} 2^0 + 2^1 + \dots + 2^k + 2^{k+1} &= (2^0 + 2^1 + \dots + 2^k) + 2^{k+1} \\ &= (2^{k+1} - 1) + 2^{k+1} && \text{by the IH} \\ &= 2(2^{k+1}) - 1 \\ &= 2^{k+2} - 1, \end{aligned}$$

which is exactly $P(k+1)$.

5. Thus $P(n)$ is true for all $n \in \mathbb{N}$, by induction.
- 

Prove $1 + 2 + 3 + \dots + n = n(n + 1)/2$

Prove $1 + 2 + 3 + \dots + n = n(n + 1)/2$

1. Let $P(n)$ be “ $0 + 1 + 2 + \dots + n = n(n+1)/2$ ”. We will show $P(n)$ is true for all natural numbers by induction.

Summation Notation

$$\sum_{i=0}^n i = 0 + 1 + 2 + 3 + \dots + n$$

Prove $1 + 2 + 3 + \dots + n = n(n + 1)/2$

- 1. Let $P(n)$ be “ $0 + 1 + 2 + \dots + n = n(n+1)/2$ ”. We will show $P(n)$ is true for all natural numbers by induction.**
- 2. Base Case ($n=0$): $0 = 0(0+1)/2$. Therefore $P(0)$ is true.**

Summation Notation

$$\sum_{i=0}^n i = 0 + 1 + 2 + 3 + \dots + n$$

Prove $1 + 2 + 3 + \dots + n = n(n + 1)/2$

1. Let $P(n)$ be " $0 + 1 + 2 + \dots + n = n(n+1)/2$ ". We will show $P(n)$ is true for all natural numbers by induction.
2. Base Case ($n=0$): $0 = 0(0+1)/2$. Therefore $P(0)$ is true.
3. Induction Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 0$. I.e., suppose $1 + 2 + \dots + k = k(k+1)/2$

"some" or "an"
not any!

Prove $1 + 2 + 3 + \dots + n = n(n + 1)/2$

1. Let $P(n)$ be " $0 + 1 + 2 + \dots + n = n(n+1)/2$ ". We will show $P(n)$ is true for all natural numbers by induction.
2. Base Case ($n=0$): $0 = 0(0+1)/2$. Therefore $P(0)$ is true.
3. Induction Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 0$. I.e., suppose $1 + 2 + \dots + k = k(k+1)/2$

Prove $1 + 2 + 3 + \dots + n = n(n + 1)/2$

1. Let $P(n)$ be " $0 + 1 + 2 + \dots + n = n(n+1)/2$ ". We will show $P(n)$ is true for all natural numbers by induction.
2. Base Case ($n=0$): $0 = 0(0+1)/2$. Therefore $P(0)$ is true.
3. Induction Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 0$. I.e., suppose $1 + 2 + \dots + k = k(k+1)/2$
4. Induction Step:
→ Goal: Show $P(k+1)$, i.e. show $1 + 2 + \dots + k + (k+1) = (k+1)(k+2)/2$

Prove $1 + 2 + 3 + \dots + n = n(n + 1)/2$

1. Let $P(n)$ be " $0 + 1 + 2 + \dots + n = n(n+1)/2$ ". We will show $P(n)$ is true for all natural numbers by induction.
2. Base Case ($n=0$): $0 = 0(0+1)/2$. Therefore $P(0)$ is true.
3. Induction Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 0$. I.e., suppose $1 + 2 + \dots + k = k(k+1)/2$
4. Induction Step:

$$\begin{aligned} 1 + 2 + \dots + k + (k+1) &= (1 + 2 + \dots + k) + (k+1) \\ &= k(k+1)/2 + (k+1) \text{ by IH} \\ &= (k+1)(k/2 + 1) \\ &= (k+1)(k+2)/2 \end{aligned}$$

$k/2 + 1 = \frac{k+2}{2}$

So, we have shown $1 + 2 + \dots + k + (k+1) = (k+1)(k+2)/2$, which is exactly $P(k+1)$.

5. Thus $P(n)$ is true for all $n \in \mathbb{N}$, by induction.

Induction: Changing the start line

- What if we want to prove that $P(n)$ is true for all integers $n \geq b$ for some integer b ?
- Define predicate $Q(k) = P(k + b)$ for all k .
 - Then $\forall n Q(n) \equiv \forall n \geq b P(n)$
- Ordinary induction for Q :
 - Prove $Q(0) \equiv P(b)$
 - Prove
$$\forall k (Q(k) \rightarrow Q(k + 1)) \equiv \forall k \geq b (P(k) \rightarrow P(k + 1))$$

Inductive Proofs starting at b in 5 Easy Steps

1. “Let $P(n)$ be... . We will show that $P(n)$ is true for all integers $n \geq b$ by induction.”
2. “Base Case:” Prove $P(b)$
3. “Inductive Hypothesis:
Assume $P(k)$ is true for an arbitrary integer $k \geq b$ ”
4. “Inductive Step:” Prove that $P(k + 1)$ is true:
Use the goal to figure out what you need.
Make sure you are using I.H. and point out where you are using it. (Don't assume $P(k + 1)$!!)
5. “Conclusion: $P(n)$ is true for all integers $n \geq b$ ”

Prove $3^n \geq n^2 + 3$ for all $n \geq 2$

Prove $3^n \geq n^2 + 3$ for all $n \geq 2$

- 1. Let $P(n)$ be " $3^n \geq n^2 + 3$ ". We will show $P(n)$ is true for all integers $n \geq 2$ by induction.**

Prove $3^n \geq n^2 + 3$ for all $n \geq 2$

- 1. Let $P(n)$ be “ $3^n \geq n^2 + 3$ ”. We will show $P(n)$ is true for all integers $n \geq 2$ by induction.**
- 2. Base Case ($n=2$): $3^2 = 9 \geq 7 = 4 + 3 = 2^2 + 3$ so $P(2)$ is true.**

Prove $3^n \geq n^2 + 3$ for all $n \geq 2$

1. Let $P(n)$ be " $3^n \geq n^2 + 3$ ". We will show $P(n)$ is true for all integers $n \geq 2$ by induction.
2. Base Case ($n=2$): $3^2 = 9 \geq 7 = 4 + 3 = 2^2 + 3$ so $P(2)$ is true.
3. Inductive Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 2$. I.e., suppose $3^k \geq k^2 + 3$.



Prove $3^n \geq n^2 + 3$ for all $n \geq 2$

1. Let $P(n)$ be “ $3^n \geq n^2 + 3$ ”. We will show $P(n)$ is true for all integers $n \geq 2$ by induction.
2. Base Case ($n=2$): $3^2 = 9 \geq 7 = 4 + 3 = 2^2 + 3$ so $P(2)$ is true.
3. Inductive Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 2$. I.e., suppose $3^k \geq k^2 + 3$.
4. Inductive Step:

Goal: Show $P(k+1)$, i.e. show $3^{k+1} \geq (k+1)^2 + 3$

Prove $3^n \geq n^2 + 3$ for all $n \geq 2$

1. Let $P(n)$ be “ $3^n \geq n^2 + 3$ ”. We will show $P(n)$ is true for all integers $n \geq 2$ by induction.
2. Base Case ($n=2$): $3^2 = 9 \geq 7 = 4 + 3 = 2^2 + 3$ so $P(2)$ is true.
3. Inductive Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 2$. I.e., suppose $3^k \geq k^2 + 3$.
4. Inductive Step:

Goal: Show $P(k+1)$, i.e. show $3^{k+1} \geq (k+1)^2 + 3 = k^2 + 2k + 4$

Prove $3^n \geq n^2 + 3$ for all $n \geq 2$

1. Let $P(n)$ be “ $3^n \geq n^2 + 3$ ”. We will show $P(n)$ is true for all integers $n \geq 2$ by induction.
2. Base Case ($n=2$): $3^2 = 9 \geq 7 = 4 + 3 = 2^2 + 3$ so $P(2)$ is true.
3. Inductive Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 2$. I.e., suppose $3^k \geq k^2 + 3$.
4. Inductive Step:

Goal: Show $P(k+1)$, i.e. show $3^{k+1} \geq (k+1)^2 + 3 = k^2 + 2k + 4$

$$\begin{aligned} 3^{k+1} &= 3(3^k) \\ &\geq 3(k^2 + 3) \text{ by the IH} \\ &= 3k^2 + 9 \\ &= k^2 + 2k^2 + 9 \\ &\geq k^2 + 2k + 4 = (k+1)^2 + 3 \text{ since } k \geq 2. \end{aligned}$$

Therefore $P(k+1)$ is true.

Prove $3^n \geq n^2 + 3$ for all $n \geq 2$

1. Let $P(n)$ be " $3^n \geq n^2 + 3$ ". We will show $P(n)$ is true for all integers $n \geq 2$ by induction.
2. Base Case ($n=2$): $3^2 = 9 \geq 7 = 4 + 3 = 2^2 + 3$ so $P(2)$ is true.
3. Inductive Hypothesis: Suppose that $P(k)$ is true for some arbitrary integer $k \geq 2$. I.e., suppose $3^k \geq k^2 + 3$.

4. Inductive Step:

Goal: Show $P(k+1)$, i.e. show $3^{k+1} \geq (k+1)^2 + 3 = k^2 + 2k + 4$

$$3^{k+1} = 3(3^k)$$

$$\geq 3(k^2 + 3) \text{ by the IH}$$

$$= k^2 + 2k^2 + 9$$




$$\geq k^2 + 2k + 4 = (k+1)^2 + 3 \text{ since } k \geq 2.$$

Therefore $P(k+1)$ is true.

5. Thus $P(n)$ is true for all integers $n \geq 2$, by induction.

Checkerboard Tiling

- Prove that a $2^n \times 2^n$ checkerboard with one square removed can be tiled with:



Checkerboard Tiling

1. Let $P(n)$ be any $2^n \times 2^n$ checkerboard with one square removed can be tiled with  .

We prove $P(n)$ for all $n \geq 1$ by induction on n .

Checkerboard Tiling

1. Let $P(n)$ be any $2^n \times 2^n$ checkerboard with one square removed can be tiled with  .

We prove $P(n)$ for all $n \geq 1$ by induction on n .

2. Base Case: $n=1$    

Checkerboard Tiling

1. Let $P(n)$ be any $2^n \times 2^n$ checkerboard with one square removed can be tiled with  .

We prove $P(n)$ for all $n \geq 1$ by induction on n .

2. Base Case: $n=1$    

3. Inductive Hypothesis: Assume $P(k)$ for some arbitrary integer $k \geq 1$

Checkerboard Tiling


1. Let $P(n)$ be any $2^n \times 2^n$ checkerboard with one square removed can be tiled with  .

We prove $P(n)$ for all $n \geq 1$ by induction on n .

2. Base Case: $n=1$    

3. Inductive Hypothesis: Assume $P(k)$ for some arbitrary integer $k \geq 1$

4. Inductive Step: Prove $P(k+1)$



Apply IH to each quadrant then fill with extra tile.